

Adipose-derived Stem Cells Attenuates Diabetic Osteoarthritis via Inhibition of Glycation-mediated Inflammatory Cascade

Navneet Kumar Dubey^{1,2}, Hong-Jian Wei^{2,3,4}, Sung-Hsun Yu², David F. Williams⁵, Joseph R. Wang⁶, Yue-Hua Deng⁷, Feng-Chou Tsai⁸, Peter D. Wang^{4,9}, Win-Ping Deng^{2,4,10*}

¹Graduate Institute of Biomedical Materials and Tissue Engineering, College of Biomedical Engineering, Taipei Medical University, Taipei, Taiwan

²Stem Cell Research Center, College of Oral Medicine, Taipei Medical University, Taipei, Taiwan

³School of Dental Technology, College of Oral Medicine, Taipei Medical University, Taipei, Taiwan

⁴School of Dentistry, College of Oral Medicine, Taipei Medical University, Taipei, Taiwan

⁵Wake Forest Institute of Regenerative Medicine, Winston-Salem, NC, USA

⁶Department of Periodontics, College of Dental Medicine, Columbia University, New York, USA


⁷Department of Life Science, Fu Jen Catholic University, New Taipei City, Taiwan

⁸Stem Cell Research Center, Cosmetic Clinic Group, Taipei, Taiwan

⁹Department of Dentistry, Taipei Medical University Hospital, Taipei, Taiwan

¹⁰Graduate Institute of Basic Medicine, Fu Jen Catholic University, New Taipei City, Taiwan

SUPPLEMENTARY DATA



Supplementary Figure 1. ADSC administration and evaluation of body weight in knee-joint of diabetic mice. (Control, n = 5; DM-OA+PBS, n=6; and DM-OA+ADSC, n=6). *p < 0.05 and ** p < 0.01.