

Six Visual Rating Scales as A Biomarker for Monitoring Atrophied Brain Volume in Parkinson's Disease

Yu Lin^{1,#}, Ying Fu^{1,2,#}, Yi-Fang Zeng^{1,#}, Jian-Ping Hu³, Xiao-Zhen Lin⁴, Nai-Qing Cai¹, Qiang Weng³, Yi-Jing Zhao³, Yi Lin¹, Dai-Rong Cao^{3*}, Ning Wang^{1*}

¹Department of Neurology and Institute of Neurology, The First Affiliated Hospital, Fujian Medical University, Fuzhou 350005, China.

²Central Laboratory, The First Affiliated Hospital, Fujian Medical University, Fuzhou 350005, China.

³Department of Radiology, First Affiliated Hospital, Fujian Medical University, Fuzhou 350005, China.

⁴Department of Geriatrics, First Affiliated Hospital, Fujian Medical University, Fuzhou 350005, China.

SUPPLEMENTARY DATA

Supplementary Table 1. Assessment of consistency of rating between raters.

Visual rating scales	Intraclass correlation coefficient (ICC)
Total visual rating scales	0.94
OF_L	0.81
OF_R	0.71
AC_L	0.76
AC_R	0.87
AT_L	0.81
AT_R	0.72
FI_L	0.82
FI_R	0.72
MT_L	0.76
MT_R	0.71
PA_L	0.82
PA_R	0.77

L = Left; R = Right; OF = orbitofrontal cortex; AC = anterior cingulate; FI = frontoinsula; AT = anterior temporal; MT= medial temporal lobe; PA = posterior cortex.