


Hydrogen Peroxide-Induced Senescence Reduces the Wound Healing-Promoting Effects of Mesenchymal Stem Cell-Derived Exosomes Partially via miR-146a

**Meiqian Xu^{1,#}, Xiaodong Su^{2,#}, Xian Xiao¹, Hongliang Yu¹, Xiaoxia Li³, Armand Keating^{4,5,6},
Shihua Wang^{1*}, Robert Chunhua Zhao^{1*}**

¹Institute of Basic Medical Sciences Chinese Academy of Medical Sciences, School of Basic Medicine Peking Union Medical College, Center of Excellence in Tissue Engineering Chinese Academy of Medical Sciences, Beijing 100005, China. ²Brain Tumor Research Center, Beijing Neurosurgical Institute, Beijing Tiantan Hospital Affiliated to Capital Medical University, Beijing Laboratory of Biomedical Materials, Beijing 100070, China. ³Department of Genetics and Cell Biology, Basic Medical College, Qingdao University, Qingdao 266071, China. ⁴Cell Therapy Translational Research Laboratory, Princess Margaret, Cancer Centre, Toronto, Ontario M5G 2M9, Canada. ⁵Institute of Biomaterials and Biomedical Engineering, University of Toronto, Toronto, Ontario M5G2M9, Canada. ⁶Institute of Medical Science, University of Toronto, Toronto, Ontario M5G 2M9, Canada.

SUPPLEMENTARY DATA



Supplementary Figure 1. Reduced tube numbers around wound in skins treated with s-exo. (A): Immunohistochemical stain with CD31 and ERG in skins. (B): Quantification of positive tube numbers.